


LP-2016



Publication number:

0 295 109
A1

EUROPEAN PATENT APPLICATION

Application number: 88305293.8

Int. Cl.⁴: C 07 C 69/675
C 07 C 67/317

Date of filing: 09.06.88

Priority: 11.06.87 JP 145975/87

Date of publication of application:
14.12.88 Bulletin 88/50

Designated Contracting States:
CH DE FR GB LI NL

Applicant: Takasago International Corporation
19-22, Takanawa 3-chome Minato-ku
Tokyo (JP)

Inventor: Sayo, Noboru
1151-127, Kamigo-cho Sakae-ku
Yokohama-shi Kanagawa (JP)

Akutagawa, Susumu
1080-22, Shinohara-cho Kohoku-ku
Yokohama-shi Kanagawa (JP)

Saito, Takao
298-46, Setogaya-cho Hodogaya-ku
Yokohama-shi Kanagawa (JP)

Noyori, Ryoji
135-417, Aza Shinden Ooaza Umemori Nisshin-cho
Aichi-gun Aichi (JP)


Kumobayashi, Hidenori
1-4-39, Nakakaigan
Chigasaki-shi Kanagawa (JP)

Takaya, Hidemasa
11-72, Aza Sakashita Myoudaiji-cho
Okazaki-shi Aichi (JP)

Representative: Diamond, Bryan Clive et al
Gee & Co. Chancery House Chancery Lane
London WC2A 1QU (GB)

Process for preparing optically active alcohol.

A β -keto acid derivative (1 mol) of formula (II):



kg/cm² for 1-48 hours and there is recovered as product an optically active alcohol of the formula (I) wherein the =C=O group has become CH-OH.

where R¹ is optionally substituted C1-7 alkyl, trifluoromethyl or aryl, R² is C1-8 alkoxy, SR⁵ where R⁵ is C1-8 alkyl or phenyl or -NR⁶R⁷ where R⁶ and R⁷ are H, C1-8 alkyl or benzyl, and R³ is H, halogen, C1-8 alkyl or alkoxycarbonyl of R¹+R³ form a methylene chain, is dissolved in a solvent and there is added 100-1/50,000 mol of a ruthenium-optically active phosphine derivative as catalyst, e.g. of formula Ru₂H₂Cl₂(R⁸-BINAP)₂(S)_p (III) or [RuH₂(R⁸-BINAP)₂Y_w] (V) wherein BINAP is a specified tertiary phosphine group.

The derivative is reacted with hydrogen at a pressure of 5-100

Description

PROCESS FOR PREPARING OPTICALLY ACTIVE ALCOHOL

This invention relates to a process for preparing an optically active alcohol useful as intermediate for synthesizing pharmaceuticals, liquid crystal material, and the like by asymmetric hydrogenation of a β -keto acid derivative in the presence of a ruthenium-optically active phosphine complex as a catalyst.

Known techniques for asymmetrically synthesizing optically active alcohols include a process comprising asymmetric hydrogenation using baker's yeast and a process comprising asymmetric hydrogenation using a specific catalyst.

In particular, with respect to asymmetric hydrogenation of β -keto acid derivatives to obtain optically active alcohols, it has been reported that the asymmetric hydrogenation can be carried out by using a rhodium-optically active phosphine complex as a catalyst. For example, J. Solodar reports in *Chemtech.*, 421-423 (1975) that asymmetric hydrogenation of methyl acetoacetate gives methyl 3-hydroxybutyrate in an optical yield of 71%ee.

Further, asymmetric hydrogenation using a tartaric acid-modified nickel catalyst has been proposed. According to this technique, asymmetric hydrogenation of methyl acetoacetate gives methyl 3-hydroxybutyrate in an optical yield of 85%ee as disclosed in Tai, *Yukagaku*, 822-831 (1980).


Although the process using baker's yeast produces an alcohol having relatively high optical purity, the resulting optically active alcohol is limited in absolute configuration, and synthesis of an enantiomer is difficult.

The process utilizing asymmetric hydrogenation of β -keto acid derivative in the presence of a rhodium-optically active phosphine complex does not produce an alcohol having sufficient optical purity. Besides, metallic rhodium to be used in the catalyst is expensive due to limitations in place and quantity of production. When used as a catalyst component, it forms a large proportion in cost of the catalyst, ultimately resulting in increase in cost of the final commercial products.


The process using a tartaric acid-modified nickel catalyst involves the disadvantages of difficulty in preparing the catalyst and insufficient optical yield.

As a result of extensive investigations with the purpose of meeting the above-described problems, the inventors have found that an optically active alcohol having high optical purity can be obtained by asymmetric hydrogenation of a β -keto acid derivative in the presence of a relatively cheap ruthenium-optically active phosphine complex as a catalyst. The present invention has been completed based on this finding.

The present invention relates to a process for preparing an optically active alcohol represented by formula (I):



wherein R^1 represents a substituted or unsubstituted lower alkyl group, a trifluoromethyl group or an aryl group; R^2 represents OR^4 , wherein R^4 represents an alkyl group having from 1 to 8 carbon atoms, SR^5 , wherein R^5 represents a lower alkyl group or a phenyl group, or NR^6R^7 , wherein R^6 and R^7 , which may be the same or different, each represents a hydrogen atom, a lower alkyl group or a benzyl group; and R^3 represents a hydrogen atom, a halogen atom, a lower alkyl group, a lower alkoxy carbonyl group or a lower alkoxy carbonyl-lower alkyl group; or R^1 and R^3 are connected to each other to form a methylene chain, forming a 4- to 6-membered ring together with the carbon atoms therebetween, which comprises asymmetrically hydrogenating a β -keto acid derivative represented by formula (II):



wherein R^1 , R^2 , and R^3 are as defined above,
in the presence of a ruthenium-optically active phosphine complex as a catalyst.

The "lower" alkyl or alkoxy groups have 1 to 7, preferably 1 to 4 carbon atoms.


in formulae (I) and (II), substituents for the lower alkyl group as represented by R^1 include a halogen atom, a hydroxyl group, an amino group, a lower alkyl-substituted amino group, a benzyloxy group, and an aryl group.

The β -keto acid derivative represented by formula (II) which can be used in the present invention as a starting compound specifically includes methyl acetoacetate, ethyl acetoacetate, isopropyl acetoacetate, n-butyl acetoacetate, t-butyl acetoacetate, n-pentyl acetoacetate, n-hexyl acetoacetate, n-heptyl acetoacetate, n-octyl acetoacetate, methyl 4-chloroacetoacetate, ethyl 4-chloroacetoacetate, methyl 4-fluoroacetoacetate, methyl 3-oxopentanoate, methyl 3-oxohexanoate, methyl 3-oxoheptanoate, ethyl 3-oxooctanoate, ethyl 3-oxononanoate, ethyl 3-oxodecanoate, ethyl 3-oxoundecanoate, ethyl 3-oxo-3-phenylpropanoate, ethyl 3-oxo-3-p-methoxyphenylpropanoate, ethyl 4-phenyl-3-oxobutanoate, methyl 5-phenyl-3-oxopentanoate, ethyl 3-trifluoromethyl-3-oxopropanoate, ethyl 4-hydroxy-3-oxobutanoate, methyl 4-benzyloxy-3-oxobutanoate, ethyl 4-benzyloxy-3-oxobutanoate, methyl 4-amino-3-oxobutanoate, ethyl 4-methylamino-3-oxobutanoate, methyl 4-dimethylamino-3-oxobutanoate, ethyl 4-dimethylamino-3-oxobutanoate, ethyl 2-methylacetoacetate, ethyl 2-chloroacetoacetate, diethyl 2-acetylsuccinate, diethyl 2-acetylglutamate, 2-carboethoxy-cyclopentanone, 2-carboethoxy-cyclohexanone, dimethyl acetylmalonate, 3-oxobutanoic dimethylamide, 3-oxobutanoic benzylamide, thiomethyl acetoacetate, thioethyl acetoacetate and thiophenyl acetoacetate.

The ruthenium-optically active phosphine complex to be used as a catalyst include those represented by the following formulae (III) and (V):



wherein R^8 -BINAP represents a tertiary phosphine represented by formula (IV):



(IV)

wherein R^8 represents a hydrogen atom, a methyl group or a t-butyl group; S represents a tertiary amine; when y represents 0, then x represents 2, z represents 4, and p represents 1; and when y represents 1, then x represents 1, z represents 1, and p represents 0.




wherein R^8 -BINAP is as defined above; Y represents ClO_4 , BF_4 or PF_6 ; when ℓ represents 0, then v represents 1, and w represents 2; and when ℓ represents 1, then v represents 2 and w represents 1.

In formulae (III) and (V), "BINAP" represents a 2,3-bis(diphenylphosphino)-1,1'-binaphthyl moiety (hereinafter the same).


The compound of formula (III) can be obtained by the process disclosed in T. Ikariya et al., J. Chem. Soc., Chem. Commun., 922-924 (1985) and Japanese Patent Application (OPI) No. 63690/86 (the term "OPI" as used herein means "unexamined published Japanese patent application"). More specifically, the complex of formula (III) wherein y is 0 can be prepared by reacting 1 mol of $[RuCl_2(COD)]_n$ (wherein COD represents cycloocta-1,5-diene, hereinafter the same), which is obtainable by reacting ruthenium chloride and COD in an ethanol solution, and 1.2 mols of a 2,2'-bis(di-p- R^8 -phenyl phosphino)-1,1'-binaphthyl (R^8 -BINAP) in a solvent, e.g., toluene or ethanol, in the presence of 4 mols of a tertiary amine, e.g., triethylamine. The complex of formula (III) wherein y is 1 can be obtained by reacting 1 mol of $[RuCl_2(COD)]_n$, 2.25 mols of R^8 -BINAP, and 4.5 mols of a tertiary amine.

The complex of formula (V) wherein ℓ is 0, v is 1 and w is 2 can be prepared by reacting $Ru_2Cl_4(R^8-BINAP)_2(NEt_3)$ (wherein Et represents an ethyl group, hereinafter the same), which is obtained by the above-described process, with a salt represented by formula (VI):



wherein M represents Na, K, Li, Mg or Ag; and Y is as defined above,

in a solvent system comprising water and methylene chloride in the presence of a quaternary ammonium salt or quaternary phosphonium salt represented by formula (VII):



wherein R^9 , R^{10} , R^{11} , and R^{12} each represents an alkyl group having from 1 to 16 carbon atoms, a phenyl

group or a benzyl group; A represents a nitrogen atom or a phosphorus atom; and B represents a halogen atom, as a phase transfer catalyst. The reaction can be carried out by adding the reactants and the phase transfer catalyst of formula (VII) to a mixed solvent of water and methylene chloride and stirring the system. The amounts of the salt of formula (VI) and the phase transfer catalyst of formula (VII) to be added range from 2 to 10 mols, and preferably 5 mols, and from 1/100 to 1/10 mol, respectively, per mol of ruthenium. The reaction sufficiently proceeds by stirring at a temperature of from 5 to 30°C for a period of from 6 to 18 hours, and usually 12 hours. Examples of the phase transfer catalyst of formula (VII) are described in literature, i.e., W.P. Weber and G.W. Gokel, *Sokan Ido Shokubai* (Japanese translation), 1st Ed., Kagaku Dojinsha (1978). After completion of the reaction, the reaction mixture is allowed to stand still, followed by liquid separation. After the aqueous layer is removed, the methylene chloride solution is washed with water, and methylene chloride is removed by distillation under reduced pressure to obtain the desired compound.

The complex of formula (V) where ℓ is 1, v is 2 and w is 1 can be prepared by reacting $\text{RuHCl}(\text{R}^8\text{-BINAP})_2$ obtainable by the process disclosed in Japanese Patent Application (OPI) No. 63690/86 with the salt of formula (VI) in a mixed solvent of water and an organic solvent, e.g., methylene chloride, in the presence of the phase transfer catalyst of formula (VIII). The amounts of the salt of formula (VI) and the phase transfer catalyst of formula (VII) range from 2 to 10 mols, and preferably 5 mols, and from 1/100 to 1/10 mol, respectively, per mol of ruthenium. This reaction sufficiently proceeds by stirring at a temperature of from 5 to 30°C for a period of from 6 to 18 hours, and usually 12 hours.

Specific examples of the above-described ruthenium-phosphine complex according to the present invention are shown below.

$\text{Ru}_2\text{Cl}_4(\text{BINAP})_2(\text{NEt}_3)$
 $\text{Ru}_2\text{Cl}_4(\text{T-BINAP})_2(\text{NEt}_3)$
 [T-BINAP represents 2,2'-bis(di-p-tolylphosphino)-1,1'-binaphthyl]
 $\text{Ru}_2\text{Cl}_4(\text{t-Bu-BINAP})_2(\text{NEt}_3)$
 [t-Bu-BINAP represents 2,2'-bis(di-p-t-butylphenylphosphino)-1,1'-binaphthyl]
 $\text{RuHCl}(\text{BINAP})_2$
 $\text{RuHCl}(\text{T-BINAP})_2$
 $\text{RuHCl}(\text{t-Bu-BINAP})_2$
 $[\text{Ru}(\text{BINAP})](\text{ClO}_4)_2$
 $[\text{Ru}(\text{T-BINAP})](\text{ClO}_4)_2$
 $[\text{Ru}(\text{t-Bu-BINAP})](\text{ClO}_4)_2$
 $[\text{Ru}(\text{BINAP})](\text{BF}_4)_2$
 $[\text{Ru}(\text{T-BINAP})](\text{BF}_4)_2$
 $[\text{Ru}(\text{t-Bu-BINAP})](\text{BF}_4)_2$
 $[\text{Ru}(\text{BINAP})](\text{PF}_6)_2$
 $[\text{Ru}(\text{T-BINAP})](\text{PF}_6)_2$
 $[\text{RuH}(\text{BINAP})_2]\text{ClO}_4$
 $[\text{RuH}(\text{T-BINAP})_2]\text{ClO}_4$
 $[\text{RuH}(\text{BINAP})_2]\text{BF}_4$
 $[\text{RuH}(\text{T-BINAP})_2]\text{BF}_4$
 $[\text{RuH}(\text{BINAP})_2]\text{PF}_6$
 $[\text{RuH}(\text{T-BINAP})_2]\text{PF}_6$

In carrying out the present invention, a β -keto acid derivative of formula (II) is dissolved in an amphiprotic solvent, e.g., methanol, ethanol or methyl cellosolve, or a mixed solvent of such an amphiprotic solvent with another solvent such as tetrahydrofuran, toluene, benzene or methylene chloride. The solution is charged in an autoclave, and from 1/100 to 1/50,000 mol of a ruthenium-optically active phosphine complex is added thereto per mol of the β -keto acid derivative. The hydrogenation reaction is effected under stirring at a temperature of from 5 to 50°C, and preferably from 25 to 35°C, at a hydrogen pressure of from 5 to 100 kg/cm² for a period of from 1 to 48 hours. After completion of the reaction, the solvent is removed by distillation, and the residue is distilled under reduced pressure or subjected to silica gel column chromatography to thereby isolate the desired optically active alcohol of formula (I) in a substantially quantitative yield.

The present invention will now be illustrated in greater detail with reference to Reference Examples and Examples, but the invention is not limited thereto. In these examples, analytical instruments and conditions used for various analyses are as follows.

1) Gas Chromatography (GC):
 SHIMADZU GC-9A manufactured by Shimadzu Corporation
 Column: PEG-20M Silica Capillary, 0.25 mm in diameter and 25 m in length, manufactured by Gasukuro Kogyo Inc.
 Measurement Temperature: 100-250°C and increasing at a rate of 3°C/min.

2) High Performance Liquid Chromatography (HPLC):
 Hitachi Liquid Chromatography-655A-11 manufactured by Hitachi, Ltd.
 Column: Chemcopack Nucleosil 100-3, 4.6 mm in diameter and 300 mm in length, manufactured by Chemco Co.

Developing Solvent: Hexane:diethyl ether=7:3; flow rate: 1 ml/min
 Detector: UV Detector 655A (UV-254), manufactured by Hitachi, Ltd.

3) Optical Rotation:

Polarimeter DIP-4, manufactured by Nippon Bunko Bogyo K.K.

4) ³¹P NMR Spectrum:

JNM-GX400 (161 MHz) manufactured by JEOL Ltd.
 Chemical shift was determined by using 85% phosphoric acid as an external standard.

REFERENCE EXAMPLE 1Synthesis of Ru₂Cl₄[(+)-BINAP]₂(NEt₃)

(di[2,2'-bis(diphenylphosphino)-1,1'-binaphthyl]tetrachloro-diruthenium triethylamine):

To 100 ml of toluene were added 1 g (3.56 mmol) of [RuCl₂(COD)]_n, 2.66 g (4.27 mmol) of (+)-BINAP, and 1.5 g of triethylamine in a nitrogen atmosphere, and the mixture was heat-refluxed for 10 hours. The solvent was removed from the reaction mixture by distillation under reduced pressure, and the residual solid was dissolved in methylene chloride, followed by filtration through Celite filter aid. The filtrate was concentrated to dryness to obtain 3.7 g of the entitled compound as a deep brown solid.

Elemental Analysis for C₉₄H₇₉Cl₄NP₄Ru₂:

Calcd. (%):

Ru 11.96; C 66.85; H 4.71; P 7.33

Found (%):

Ru 11.68; C 67.62; H 4.97; P 6.94

³¹P NMR (CDCl₃) δ ppm:

51.06 (s), 51.98 (s), 53.87 (s), and 54.83 (s)

REFERENCE EXAMPLE 2Synthesis of [Ru((-)-T-BINAP)](ClO₄)₂ ([2,2'-bis(di-p-tolylphosphino)-1,1'-binaphthyl]ruthenium perchlorate):

In a 250 ml-volume Schlenk's tube was charged 0.54 g (0.3 mmol) of Ru₂Cl₄[(-)-T-BINAP]₂(NEt₃). After thorough displacement of the atmosphere with nitrogen gas, 60 ml of methylene chloride was added thereto, and then a solution of 0.73 g (6.0 mmols) of sodium perchlorate in 60 ml of water and a solution of 16 mg (0.06 mmol) of triethylbenzylammonium bromide in 3 ml of water were added to the mixture. The mixture was stirred at room temperature for 12 hours. After completion of the reaction, the reaction mixture was allowed to stand, and the aqueous layer was removed. The methylene chloride was removed from the organic layer by distillation under reduced pressure, and the residue was dried under reduced pressure to obtain 0.59 g (yield: 99.6%) of the entitled compound as a deep brown solid.

Elemental Analysis for C₄₈H₄₀Cl₂O₈P₂Ru:

Calcd. (%):

Ru 10.32; C 58.90; H 4.12; P 6.33

Found (%):

Ru 10.08; C 58.61; H 4.53; P 5.97

³¹P NMR (CDCl₃) δ ppm:

12.920 (d, J=41.1 Hz) and 61.402 (d, J=41.1 Hz)

EXAMPLE 1Synthesis of Methyl (3R)-(-)-3-Hydroxybutyrate

In a 200 ml-volume stainless steel-made autoclave whose atmosphere had been replaced with nitrogen were charged 10 ml (93 mmols) of methyl acetoacetate, 50 ml of methanol, and 0.5 ml of water, and 42 mg (0.025 mmol) of Ru₂Cl₄[(+)-BINAP]₂(NEt₃) as prepared in Reference Example 1 was added thereto to effect hydrogenation at a temperature of 30°C under a hydrogen pressure of 40 kg/cm² for 20 hours. The solvent was removed by distillation, and the residue was distilled under reduced pressure to obtain 10.8 g (98%) of the entitled compound having a boiling point of 72°C/17 mmHg.

The product was found to have a purity of 99.0% by GC and an optical rotation [α]_D²⁰ of -24.17° (neat).

Thirty milligrams of the resulting alcohol was esterified with (+)-α-methoxy-α-trifluoromethylphenylacetyl

chloride, and the ester was analyzed by GC and HPLC. The results revealed that the product was a mixture comprising 99.55% of methyl (3R)-(-)-3-hydroxybutyrate and 0.45% of methyl (3S)-(+)-3-hydroxybutyrate. Accordingly, the optical yield of the methyl (3R)-(-)-3-hydroxybutyrate was found to be 99.1%.

5

EXAMPLES 2 to 17

The same procedure of Example 1 was repeated, except for altering the reaction substrate, catalyst and reaction conditions as shown Table 1 below. The analytical results obtained are shown in Table 2.

10 In Examples 7, 8, 14, and 15, the optically active alcohol produced contains two asymmetric centers forming diastereomers. A ratio of the *syn* form to the *anti* form in each case was determined by HPLC, and the optical yield of each form was determined. The results obtained are separately shown in Table 3.

15

20

25

30

35

40

45


50

55

60


65

TABLE 1




Example No.	R ¹	R ²	R ³	Catalyst	Substrate/Catalyst (mol/mol)	Hydrogen Pressure (kg/cm ²)	Temperature (°C)	Time (hr)
2	CH ₃	OC ₂ H ₅	H	Ru ₂ Cl ₄ [(+)-BINAP] ₂ (NEt ₃)	2000	40	30	22
3	CH ₃	OlPr	H	[Ru((-)-BINAP)](ClO ₄) ₂	1000	5	30	20
4	CH ₃	OlBu	H	[Ru((-)-T-BINAP)](BF ₄) ₂	1000	5	30	20
5	CH ₃ CH ₂	OCH ₃	H	Ru ₂ Cl ₄ [(+)-BINAP] ₂ (NEt ₃)	2000	40	30	18
6	CH ₃ (CH ₂) ₃	OCH ₃	H	Ru ₂ Cl ₄ [(+)-T-BINAP] ₂ (NEt ₃)	1000	40	30	18
7	CH ₃	OC ₂ H ₅	Cl	[RuH((+)-BINAP) ₂]ClO ₄	1000	30	30	15
8	CH ₃	OC ₂ H ₅	CH ₃	Ru ₂ Cl ₄ [(+)-T-BINAP] ₂ (NEt ₃)	1000	80	30	24
9	CF ₃	OC ₂ H ₅	H	RuHCl[(+)-BINAP] ₂	1000	80	30	16
10	PhCH ₂ OCH ₂	OCH ₃	H	Ru ₂ Cl ₄ [(+)-T-BINAP] ₂ (NEt ₃)	1000	40	30	20
11	CH ₃	NHCH ₂ Ph	H	[Ru((-)-T-BINAP)](PF ₆) ₂	1000	40	30	20
12	(CH ₃) ₂ NCH ₂	OC ₂ H ₅	H	Ru ₂ Cl ₄ [(+)-BINAP] ₂ (NEt ₃)	1000	40	30	18
13	CH ₃	SC ₂ H ₅	H	[Ru((+)-T-BINAP)](ClO ₄) ₂	1000	40	30	30

TABLE 1 (cont'd)



Example No.	R ¹	R ²	R ³	Catalyst	Substrate/ Catalyst (mol/mol)	Hydrogen Pressure (kg/cm ²)	Tempera- ture (°C)	Time (hr)
14				[Ru((+)-BINAP)](BF ₄) ₂	1000	40	30	24
15	CH ₃	OCH ₃	CH ₂ CO ₂ CH ₃	RuHCl[(-)-T-BINAP] ₂	1000	40	30	36
16	ClCH ₂	CH ₃	H	Ru ₂ Cl ₄ [(+)-BINAP] ₂ (NEt ₃)	1000	100	30	16
17	BfCH ₂	CH ₃	H	Ru ₂ Cl ₄ [(-)-BINAP] ₂ (NEt ₃)	1000	40	30	20



Note: iPr represents an isopropyl group; tBu represents a t-butyl group; and Ph represents a phenyl group.

TABLE 2










<u>Example No.</u>	<u>Product</u>	<u>Yield (%)</u>	<u>Optical Yield (%ee)</u>
2		99	99.1
3		98	98.0
4		98	96.4
5		99	99.3
6		99	99.2
7		95	see Table 3
8		97	see Table 3
9		95	46
10		97	95

TABLE 2 (cont'd)








Example No.	Product	Yield (%)	Optical Yield (%ee)
11		94	88
12		91	93
13		87	65
14		90	see Table 3
15		85	see Table 3
16		90	67
17		95	45

TABLE 3

Example No.	Syn:Anti Ratio	Optical Yield (%ee)	
		Syn	Anti
7	60:40	92	88
8	50:50	90	87
14	55:45	91	89
15	60:40	92	86

As described above, the present invention provides an industrially valuable process for preparing a useful optically active alcohol at high efficiency.

Claims

1. A process for preparing an optically active alcohol represented by formula (I):



wherein R^1 represents a substituted or unsubstituted alkyl group having 1 to 7 carbon atoms, a trifluoromethyl group or an aryl group; R^2 represents OR^4 , wherein R^4 represents an alkyl group having from 1 to 8 carbon atoms, SR^5 , wherein R^5 represents an alkyl group having 1 to 7 carbon atoms or a phenyl group, or NR^6R^7 , wherein R^6 and R^7 , which may be the same or different, each represents a hydrogen atom, an alkyl group having 1 to 7 carbon atoms or a benzyl group; and R^3 represents a hydrogen atom, a halogen atom, an alkyl group having 1 to 7 carbon atoms, a C1-8 alkoxy carbonyl group or a C1-8 alkoxy carbonyl-C1-7 alkyl group; or R^1 and R^3 are connected to each other to form a methylene chain, forming a 4- to 6-membered ring together with the carbon atoms therebetween, which comprises asymmetrically hydrogenating a β -keto acid derivative represented by formula (II):




wherein R^1 , R^2 , and R^3 are as defined above, in the presence of a ruthenium-optically active phosphine complex as a catalyst.

2. A process as claimed in Claim 1, wherein said ruthenium-optically active phosphine complex is a compound represented by formula (III):



wherein $\text{R}^8\text{-BINAP}$ represents a tertiary phosphine represented by formula (IV):



wherein R^8 represents a hydrogen atom, a methyl group or a *t*-butyl group; S represents a tertiary amine; when y represents 0, then x represents 2, z represents 4, and p represents 1; and when y represents 1, then x represents 1, z represents 1, and p represents 0, or a compound represented by formula (V):



wherein $\text{R}^8\text{-BINAP}$ is as defined above; Y represents ClO_4 , BF_4 or PF_6 ; when ℓ represents 0, then v represents 1, and w represents 2; and when ℓ represents 1, then v represents 2 and w represents 1.

3. A process as claimed in Claim 1 or 2, wherein the derivative formulae (II) is dissolved in an amphiprotic solvent and the phosphine complex is added in an amount of 1/100 to 1/50,000 mol per mol of the derivative (I).



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 88305293.8
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. ⁴)
A	CHEMICAL ABSTRACTS, vol. 83, no. 13, September 29, 1975, Columbus, Ohio, USA J.SOLODAR "Asymmetric hydrogenation of ketones." page 488, column 2, abstract-no. 113 607k	1	C 07 C 69/675 C 07 C 67/317
D,A	& Chem.Technol. 1975, 5(7), 421-23 --		
A	CHEMICAL ABSTRACTS, vol. 94, no. 15, April 13, 1981, Columbus, Ohio, USA A.TAI "Asymmetrically modified nickel catalyst and its use for the preparation of optically active 3-hydroxy acids." page 631, column 1, abstract-no. 120 759s	1	
D,A	& Yukagaku 1980, 29(11), 822-31 --		
A	US - A - 4 273 933 (TADAO HARADA et al.) * Examples * ----	1,3	
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 09-08-1988	Examiner HOFBAUER
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			